

Tennessee 7th GRADE MATH 2021-2022 Pacing Guide CASE Benchmark Assessments

Unit	Standards	Major Topics/Concepts
Ratios and Proportions	7.RP.A.1 7.RP.A.2 7.RP.A.3	Compute unit rates associated with ratios of fractions, including ratios of lengths, areas, and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour. Recognize and represent proportional relationships between quantities. Decide whether two quantities are in a proportional relationship (e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin). Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships. Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p, the relationship between the total cost and the number of items can be expressed as t = pn. Explain what a point (x, y) on the graph of a proportional relationship means in terms of the situation, with special attention to the points (0, 0) and (1, r) where r is the unit rate. Use proportional relationships to solve multi-step ratio and percent problems. Examples: simple interest, tax, markups and markdowns, gratuities and commissions, fees, percent increase and decrease, percent error.
Operations with Rational Numbers	7.NS.A.1 7.NS.A.2 7.NS.A.3	 Apply and extend previous understandings of addition and subtraction to add and subtract rational numbers; represent addition and subtraction on a horizontal or vertical number line diagram. ✓ Describe situations in which opposite quantities combine to make 0. ✓ Understand p + q as the number located a distance q from p, in the positive or negative direction depending on whether q is positive or negative. Show that a number and its opposite have a sum of 0 (are additive inverses). Interpret sums of rational numbers by describing real-world contexts. ✓ Understand subtraction of rational numbers as adding the additive inverse, p - q = p + (-q). Show that the distance between two rational numbers on the number line is the absolute value of their difference, and apply this principle in real-world contexts.

Unit	Standards	Major Topics/Concepts
		✓ Apply properties of operations as strategies to add and
		subtract rational numbers.
		Apply and extend previous understandings of multiplication and division and of fractions to multiply and divide rational numbers.
		✓ Understand that multiplication is extended from fractions to
		rational numbers by requiring that operations continue to
		satisfy the properties of operations, particularly the Distributive
		Property, leading to products such as $(-1)(-1) = 1$ and the
		rules for multiplying signed numbers. Interpret products of
		rational numbers by describing real-world contexts. ✓ Understand that integers can be divided, provided that the
		divisor is not zero, and every quotient of integers (with non-
		zero divisor) is a rational number. If p and q are integers,
		then $-(p/q) = (-p)/q = p/(-q)$. Interpret quotients of
		rational numbers by describing real-world contexts.
		 Apply properties of operations as strategies to multiply and divide rational numbers.
		✓ Convert a rational number to a decimal using long division;
		know that the decimal form of a rational number terminates in
		Os or eventually repeats.
		Solve real-world and mathematical problems involving the four
		operations with rational numbers. (Computations with rational
		numbers extend the rules for manipulating fractions to complex
		fractions.)
	(c	1 st Cumulative Benchmark covering all content to this point)
		Apply properties of operations as strategies to add, subtract, factor,
		and expand linear expressions with rational coefficients.
		Understand that rewriting an expression in different forms in a
		contextual problem can provide multiple ways of interpreting the
		problem and how the quantities in it are related. For example, shoes
		are on sale at a 25% discount. How is the discounted price P related
		to the original cost C of the shoes? $C - 0.25C = P$. In other words, P is 75% of the original cost for $C - 0.25C$ can be written as 0.75C.
		Solve multi-step real-world and mathematical problems posed with
Expressions and	7.EE.A.1 7.EE.A.2	positive and negative rational numbers presented in any form (whole numbers, fractions, and decimals).
Equations	7.EE.A.2 7.EE.B.3	✓ Apply properties of operations to calculate with numbers in any
quations	7.EE.B.4	form; convert between forms as appropriate.
		✓ Assess the reasonableness of answers using mental
		computation and estimation strategies.
		Use variables to represent quantities in a real-world or mathematical
		problem, and construct simple equations and inequalities to solve
		problems by reasoning about the quantities. ✓ Solve contextual problems leading to equations of the form
		px + q = r and $p(x + q) = r$, where p , q , and r are
		specific rational numbers. Solve equations of these forms
		fluently. Compare an algebraic solution to an arithmetic
		solution, identifying the sequence of the operations used in

Unit	Standards	Major Topics/Concepts		
Construction and Deconstruction	7.G.A.1	each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width? ✓ Solve contextual problems leading to inequalities of the form px + q > r or px + q < r, where p, q, and r are specific rational numbers. Graph the solution set of the inequality on a number line, and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions. (Note that inequalities using >, <, ≤, ≥ are included in this standard.) Solve problems involving scale drawings of geometric figures, including computing actual lengths and areas from a scale drawing and reproducing a scale drawing at a different scale.		
of Geometric Figures	7.G.A.2	Draw geometric shapes with given conditions. Focus on constructing triangles from three measures of angles or sides, noticing when the conditions determine a unique triangle, more than one triangle, or no triangle.		
2 nd Cumulative Benchmark (covering all content to this point)				
Angles, Area, and Volume	7.G.B.3 7.G.B.4 7.G.B.5	Know the formulas for the area and circumference of a circle and use them to solve problems; give an informal derivation of the relationship between the circumference and area of a circle. Know and use facts about supplementary, complementary, vertical, and adjacent angles in a multi-step problem to write and solve simple equations for an unknown angle in a figure. Solve real-world and mathematical problems involving area, volume, and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms.		
Statistics	7.SP.A.1 7.SP.A.2 7.SP.B.3 7.SP.B.4	Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences. Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be. Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, measuring the difference between the centers by expressing it as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team; on a dot plot or box plot, the separation between the two		

Unit	Standards	Major Topics/Concepts		
		distributions of heights is noticeable.		
		Use measures of center and measures of variability for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a 7th grade science book are generally longer than the words in a chapter of a 4th grade science book. Understand that the probability of a chance event is a number between 0 and 1 that expresses the likelihood of the event occurring.		
Probability	7.SP.C.5 7.SP.C.6 7.SP.C.7 7.SP.D.8	Larger numbers indicate greater likelihood. A probability near 0 indicates an unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event.		
		Approximate the probability of a chance event by collecting data on the chance process that produces it and observing its long-run relative frequency, and predict the approximate relative frequency given the probability. For example, when rolling a number cube 600 times, predict that a 3 or 6 would be rolled roughly 200 times but probably not exactly 200 times.		
		Develop a probability model, and use it to find probabilities of events. Compare probabilities from a model to observed frequencies; if the agreement is not good, explain possible sources of the discrepancy. ✓ Develop a uniform probability model by assigning equal probability to all outcomes, and use the model to determine probabilities of events. For example, if a student is selected at random from a class, find the probability that Jane will be selected and the probability that a girl will be selected. ✓ Develop a probability model (which may not be uniform) by observing frequencies in data generated from a chance process. For example, find the approximate probability that a spinning penny will land heads up or that a tossed paper cup will land open-end down. Do the outcomes for the spinning penny appear to be equally likely based on the observed frequencies?		
		Summarize numerical data sets in relation to their context. ✓ Give quantitative measures of center (median and/or mean) and variability (range and/or interquartile range), and describe any overall pattern and any striking deviations from the overall pattern with reference to the context in which the data were gathered. ✓ Know and relate the choice of measures of center (median and/or mean) and variability (range and/or interquartile range) to the shape of the data distribution and the context in which the data were gathered.		
Final Comprehensive Benchmark (covering all content)				
(covering all content)				